

Hydroxychloroquine sulphate 200 milligram (mg) tablets.

The tablets are for oral administration.

BATCH NUMBER:

PRINCIPLE trial.

Chief Investigator (Butler), University of Oxford

TRIAL PARTICIPANT ID number:

Take one tablet (200mg) hydroxychloroquine twice daily for 7 days by mouth (14 tablets in total)

Special instructions:

Each dose should be taken with a meal or glass of milk.

Antacids may reduce absorption of hydroxychloroquine so it is advised that a 4-hour interval be observed between taking hydroxychloroquine and an antacid.

For clinical trial use only

Store below 25°C

Period of use: 7 (seven) days

Shelf life is 36 months.

Keep out of reach of children

Sponsor: University of Oxford, Joint Research Office, 1st floor, Boundary Brook House

Churchill Drive, Headington Oxford, OX3 7GB Tel: +44 (0)1865572224 Fax: +44 (0)1865572228

Hydroxychloroquine sulphate 200 milligram (mg) tablets.

The tablets are for oral administration.

BATCH NUMBER:

PRINCIPLE trial.

Chief Investigator (Butler), University of Oxford

TRIAL PARTICIPANT ID number:

Take one tablet (200mg) hydroxychloroquine twice daily for 7 days by mouth (14 tablets in total)

Special instructions:

Each dose should be taken with a meal or glass of milk.

Antacids may reduce absorption of hydroxychloroquine so it is advised that a 4-hour interval be observed between taking hydroxychloroquine and an antacid.

For clinical trial use only

Store below 25°C

Period of use: 7 (seven) days

Shelf life is 36 months.

Keep out of reach of children

Sponsor: University of Oxford, Joint Research Office, 1st floor, Boundary Brook House

Churchill Drive, Headington Oxford, OX3 7GB Tel: +44 (0)1865572224 Fax: +44 (0)1865572228

Hydroxychloroquine sulphate 200 milligram (mg) tablets.

The tablets are for oral administration.

BATCH NUMBER:

PRINCIPLE trial.

Chief Investigator (Butler), University of Oxford

TRIAL PARTICIPANT ID number:

Take one tablet (200mg) hydroxychloroquine twice daily for 7 days by mouth (14 tablets in total)

Special instructions:

Each dose should be taken with a meal or glass of milk.

Antacids may reduce absorption of hydroxychloroquine so it is advised that a 4-hour interval be observed between taking hydroxychloroquine and an antacid.

For clinical trial use only

Store below 25°C

Period of use: 7 (seven) days

Shelf life is 36 months.

Keep out of reach of children

Sponsor: University of Oxford, Joint Research Office, 1st floor, Boundary Brook House

Churchill Drive, Headington Oxford, OX3 7GB Tel: +44 (0)1865572224 Fax: +44 (0)1865572228

Hydroxychloroquine sulphate 200 milligram (mg) tablets.

The tablets are for oral administration.

BATCH NUMBER:

PRINCIPLE trial.

Chief Investigator (Butler), University of Oxford

TRIAL PARTICIPANT ID number:

Take one tablet (200mg) hydroxychloroquine twice daily for 7 days by mouth (14 tablets in total)

Special instructions:

Each dose should be taken with a meal or glass of milk.

Antacids may reduce absorption of hydroxychloroquine so it is advised that a 4-hour interval be observed between taking hydroxychloroquine and an antacid.

For clinical trial use only

Store below 25°C

Period of use: 7 (seven) days

Shelf life is 36 months.

Keep out of reach of children

Sponsor: University of Oxford, Joint Research Office, 1st floor, Boundary Brook House

Churchill Drive, Headington Oxford, OX3 7GB Tel: +44 (0)1865572224 Fax: +44 (0)1865572228